

**CORRECTED
VERSION***

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 38/31, 7/48		A1	(11) International Publication Number: WO 98/51331 (43) International Publication Date: 19 November 1998 (19.11.98)
(21) International Application Number: PCT/EP98/02999 (22) International Filing Date: 13 May 1998 (13.05.98) (30) Priority Data: 08/854,941 13 May 1997 (13.05.97) US (71) Applicant (for all designated States except US): SOCIETE DE CONSEILS DE RECHERCHES ET D'APPLICATIONS SCIENTIFIQUES S.A. (S.C.R.A.S.) [FR/FR]; 51, 53, rue du Docteur Blanche, F-75016 Paris (FR). (72) Inventors; and (75) Inventors/Applicants (for US only): CAWTHORNE, Michael, Anthony [GB/GB]; Clore Laboratory, University of Buckingham, Hunter Street, Buckingham, Bucks MK18 1EG (GB). LIU, Yong-Ling [GB/GB]; Clore House, Hunter Street, Buckingham, Bucks MK18 1EG (GB). SENNITT, Matthew, V. [GB/GB]; Clore House, Hunter Street, Buckingham, Bucks MK18 1EG (GB). (74) Agent: LUNT, Mark, George, Francis; Dibb Lupton Alsop, Fountain Precinct, Balm Green, Sheffield S1 1RZ (GB).			(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: SOMATOSTATIN AND SOMATOSTATIN AGONISTS FOR DECREASING BODY WEIGHT			
(57) Abstract The present invention relates to a method of decreasing body weight in a patient. The method includes the step of administering a therapeutically effective amount of a somatostatin or a somatostatin agonist to said patient. A pharmaceutical/cosmetic composition comprises the somatostatin or somatostatin agonist. Such products are used to prepare such compositions for the reduction of body weight in a human or mammalian animal.			

*(Referred to in PCT Gazette No. 12/1999, Section II)

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

SOMATOSTATIN AND SOMATOSTATIN AGONISTS FOR DECREASING BODY WEIGHT

5 This invention relates to a method and
composition useful for reducing body weight in human or
mammalian animal bodies.

BACKGROUND OF THE INVENTION

10 An estimated 35 million Americans are at least 20%
overweight (Biotechnology 13:1060-1063 (1995)), a level
at which health risks are significantly elevated. Nearly
twice this number of Americans believe themselves to be
overweight. A comparable picture is reported elsewhere.
15 For example, in the United Kingdom, approximately one
third of the women and 43% of the men are overweight,
with at least one in six women and one in eight men
classifiable as medically obese (Purnell, S., Highfield,
The Daily Telegraph, Sept. 30, 1995). There, therefore,
20 are both aesthetic and health reasons for weight control.

 In the medically obese population, the condition
is more severe and often associated with a myriad of
serious medical problems such as non-insulin dependent
diabetes mellitus, hypertension, dyslipidemia, coronary
25 heart disease and musculoskeletal disorders. Thus,
obesity is not just a problem of passive increase in
adipose mass. It has been suggested that the underlying
metabolic alterations in obesity may be amenable to
therapeutic intervention (Goldstein, D.J., et al., Am. J.
30 Clin. Nutr., 60:647-657 (1994)).

CONFIRMATION COPY

SUMMARY OF THE INVENTION

The present invention relates to a method of decreasing body weight in a patient (e.g., a mammal such as a human). The method includes the step of
5 administering a therapeutically effective amount of somatostatin or a somatostatin agonist to said patient. The somatostatin or somatostatin agonist may be administered parenterally, e.g., administered intravenously, subcutaneously, or by implantation of a
10 sustained release formulation. In one embodiment, the patient is obese (e.g., as defined by either 20-25% over normal body weight (Statistical Bulletin, Metropolitan Life Insurance Co., Vol. 40, pg. 1 (1959) or as defined by body mass index (BMI) greater than 25% over normal and
15 including risk factors or a BMI greater than 30% over normal (see, e.g., Bray, GA and Gray, DS, Diabetes/Metabolism Review 4:653-679 (1988); Flynn, et al., Proc. Nutritional Society 50:413 (1991)). In another embodiment, the patient is a non-insulin
20 dependent diabetic (i.e., type-2 diabetic).

The invention also comprises a pharmaceutical or cosmetic composition comprising a somatostatin or a somatostatin agonist. It further comprises the use of such compositions in the preparation of a pharmaceutical
25 or cosmetic composition for the reduction of excessive body weight in a human or mammalian animal.

The term "somatostatin agonist" will be defined below. A therapeutically effective amount depends upon the condition being treated, the route of administration
30 chosen, and the specific activity of the compound used

and ultimately will be decided by the attending physician or veterinarian (e.g., between 5 µg/day to 5 mg/day). In one embodiment, the somatostatin agonist is administered to the patient until the patient has lost the requisite
5 amount of body weight (e.g., the patient is no longer medically obese). In another embodiment, the somatostatin agonist is administered for the lifetime of the patient (e.g., maintaining normal body weight or secondary endpoints). In another embodiment, the
10 somatostatin agonist is administered for cosmetic purposes.

The somatostatin agonist may be injected parenterally, e.g., intravenously, into the bloodstream of the subject being treated. However, it will be
15 readily appreciated by those skilled in the art that the route, such as intravenous, subcutaneous, intramuscular, intraperitoneal, enterally, transdermally, transmucously, sustained released polymer compositions (e.g., a lactic acid polymer or copolymer microparticle or implant),
20 profusion, nasal, oral, etc., will vary with the condition being treated and the activity and bioavailability of the somatostatin agonist being used.

While it is possible for the somatostatin agonist to be administered as the pure or substantially pure
25 compound, it may also be presented as a pharmaceutical formulation or preparation. The formulations to be used in the present invention, for both humans and animals, comprise any of the somatostatin agonists to be described below, together with one or more pharmaceutically

acceptable carriers thereof, and optionally other therapeutic ingredients.

The carrier must be "acceptable" in the sense of being compatible with the active ingredient(s) of the formulation (e.g., capable of stabilizing peptides) and not deleterious to the subject to be treated. Desirably, the formulation should not include oxidizing agents or other substances with which peptides are known to be incompatible. For example, somatostatin agonists in the cyclized form (e.g., internal cysteine disulfide bond) are oxidized; thus, the presence of reducing agents as excipients could lead to an opening of the cysteine disulfide bridge. On the other hand, highly oxidative conditions can lead to the formation of cysteine sulfoxide and to the oxidation of tryptophan. Consequently, it is important to carefully select the excipient. pH is another key factor, and it may be necessary to buffer the product under slightly acidic conditions (pH 5 to 6).

The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing the active ingredient(s) into association with the carrier which constitutes one or more accessory ingredients.

In general, the formulations for tablets or powders are prepared by uniformly and intimately blending the active ingredient with finely divided solid carriers, and then, if necessary, as in the case of tablets, forming the product into the desired shape and size.

Formulations suitable for parenteral (e.g., intravenous) administration, on the other hand, conveniently comprise sterile aqueous solutions of the active ingredient(s). Preferably, the solutions are isotonic with the blood of the subject to be treated. Such formulations may be conveniently prepared by dissolving solid active ingredient(s) in water to produce an aqueous solution, and rendering said solution sterile. The formulation may be presented in unit or multi-dose containers, for example, sealed ampoules or vials.

Formulations suitable for sustained release parenteral administrations (e.g., biodegradable polymer formulations such as polyesters containing lactic or glycolic acid residues) are also well known in the art. See, e.g., U.S. Patent Nos. 3,773,919 and 4,767,628 and PCT Publication No. WO 94/15587.

The somatostatin or somatostatin agonist may also be administered with other antiobesity agents such as phentermine, diethylpropion, methamphetamine, phendimetrazine, phenmetrazine, diethylpropion, phentermine, mazindol, dextroamphetamine, phentermine, bezphetamine, orlistat, β 3-adrenergic agonists (e.g., BTA-234 and SR58611A), sibutramine, henylpropanolamine, dexfenturamine, leptin, or bromocriptine.

Other features and advantages of the invention will be apparent from the following description of the preferred embodiments and from the claims.

Abbreviations

- β -Nal = β -naphthylalanine
 β -Pal = β -pyridylalanine
5 hArg(Bu) = N-guanidino-(butyl)-homoarginine
hArg(Et)₂ = N, N'-guanidino-(dimethyl)-homoarginine
hArg(CH₂CF₃)₂ = N, N'-guanidino-bis-(2,2,2,-
trifluoroethyl) - homoarginine
hArg(CH₃, hexyl) = N, N'-guanidino-(methyl, hexyl) -
10 homoarginine
Lys(Me) = N'-methyllysine
Lys(iPr) = N'-isopropyllysine
AmPhe = aminomethylphenylalanine
AChxAla = aminocyclohexylalanine
15 Abu = α -aminobutyric acid
Tpo = 4-thiaproline
MeLeu = N-methyllleucine
Orn = ornithine
Nle = norleucine
20 Nva = norvaline
Trp(Br) = 5-bromo-tryptophan
Trp(F) = 5-fluoro-tryptophan
Trp(NO₂) = 5-nitro-tryptophan
Gaba = γ -aminobutyric acid
25 Bmp = β -mercaptopropionyl
Ac = acetyl
Pen = pencillamine

DETAILED DESCRIPTION OF THE INVENTION

- 30 It is believed that one skilled in the art can,
based on the description herein, utilize the present

invention to its fullest extent. The following specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

5 Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Also, all publications, patent applications, patents, and other references
10 mentioned herein are incorporated by reference.

Somatostatin and its Agonists

Somatostatin (somatotropin release inhibiting factor or SRIF) has both a 14 amino acid isoform
15 (somatostatin-14) and a 28 amino acid isoform (somatostatin-28). See Wilson, J. & Foster, D., *Williams Textbook of Endocrinology*, p. 510 (7th ed., 1985). The compound is an inhibitor of secretion of the growth hormone and was originally isolated from the
20 hypothalamus. Brazeau, et al., *Science* 179:77 (1973). Native somatostatin has a very short duration of effect *in vivo* since it is rapidly inactivated by endo- and exopeptidase. Many novel analogs have been prepared in order to enhance the duration of effect, biological
25 activity, and selectivity (e.g., for the particular somatostatin receptor) of this hormone. Such analogs will be called "somatostatin agonists" herein.

Various somatostatin receptors (SSTRs) have been isolated, e.g., SSTR-1, SSTR-2, SSTR-3, SSTR-4, and SSTR-
30 5. Thus, the somatostatin agonist may be a SSTR-1

agonist, SSTR-2 agonist, SSTR-3 agonist, SSTR-4 agonist or an SSTR-5 agonist. In one embodiment, the somatostatin agonist of the present invention is an SSTR-5 agonist or an SSTR-2 agonist. What is meant by an "SSTR-5 agonist" or an "SSTR-2 agonist" is a compound which (1) has a high affinity (e.g., K_i of less than 1 μM or, preferably, of less than 10 nM, or less than 2 nM, or of less than 1 nM) for the SSTR-5 or SSTR-2, respectively (e.g., as defined by the receptor binding assay described below), and (2) decreases body weight of a patient (e.g., as defined by the biological assay described below). The somatostatin agonist may also be selective for a particular somatostatin receptor, e.g., have a higher binding affinity for a particular somatostatin receptor subtype as compared to the other receptor subtypes. In one embodiment, the somatostatin receptor is an SSTR-5 selective agonist or SSTR-2 selective agonist. What is meant by an SSTR-5 selective agonist is a somatostatin agonist which (1) has a higher binding affinity (i.e., K_i) for SSTR-5 than for either SSTR-1, SSTR-2, SSTR-3, or SSTR-4 and (2) decreases body weight of a patient (e.g., as defined by the biological assay described below). In one embodiment, the SSTR-5 selective agonist has a K_i for SSTR-5 that is at least 2 times (e.g., at least 5 times or at least 10 times) less than its K_i for the SSTR-2 receptor (e.g., as defined by the receptor binding assay described below).

Somatostatin agonists which can be used to practice the therapeutic method of the present invention include, but are not limited to, those covered by

formulae or those specifically recited in the publications set forth below, all of which are hereby incorporated by reference.

- EP Application No. P5 164 EU (Inventor: G. Keri);
5 Van Binst, G. et al. Peptide Research 5:8 (1992);
Horvath, A. et al. Abstract, "Conformations of Somatostatin Analogs Having Antitumor Activity", 22nd European peptide Symposium, September 13-19, 1992, Interlaken, Switzerland;
- 10 PCT Application WO 91/09056 (1991);
EP Application 0 363 589 A2 (1990);
U.S. Patent No. 4,904,642 (1990);
U.S. Patent No. 4,871,717 (1989);
U.S. Patent No. 4,853,371 (1989);
15 U.S. Patent No. 4,725,577 (1988);
U.S. Patent No. 4,684,620 (1987)
U.S. Patent No. 4,650,787 (1987);
U.S. Patent No. 4,603,120 (1986);
U.S. Patent No. 4,585,755 (1986);
20 EP Application 0 203 031 A2 (1986);
U.S. Patent No. 4,522,813 (1985);
U.S. Patent No. 4,486,415 (1984);
U.S. Patent No. 4,485,101 (1984);
U.S. Patent No. 4,435,385 (1984);
25 U.S. Patent No. 4,395,403 (1983);
U.S. Patent No. 4,369,179 (1983);
U.S. Patent No. 4,360,516 (1982);
U.S. Patent No. 4,358,439 (1982);
U.S. Patent No. 4,328,214 (1982);
30 U.S. Patent No. 4,316,890 (1982);

10

- U.S. Patent No. 4,310,518 (1982);
U.S. Patent No. 4,291,022 (1981);
U.S. Patent No. 4,238,481 (1980);
U.S. Patent No. 4,235,886 (1980);
5 U.S. Patent No. 4,224,190 (1980);
U.S. Patent No. 4,211,693 (1980);
U.S. Patent No. 4,190,648 (1980);
U.S. Patent No. 4,146,612 (1979);
U.S. Patent No. 4,133,782 (1979);
10 U.S. Patent No. 5,506,339 (1996);
U.S. Patent No. 4,261,885 (1981);
U.S. Patent No. 4,728,638 (1988);
U.S. Patent No. 4,282,143 (1981);
U.S. Patent No. 4,215,039 (1980);
15 U.S. Patent No. 4,209,426 (1980);
U.S. Patent No. 4,190,575 (1980);
EP Patent No. 0 389 180 (1990);
EP Application No. 0 505 680 (1982);
EP Application No. 0 083 305 (1982);
20 EP Application No. 0 030 920 (1980);
PCT Application No. WO 88/05052 (1988);
PCT Application No. WO 90/12811 (1990);
PCT Application No. WO 97/01579 (1997);
PCT Application No. WO 91/18016 (1991);
25 U.K. Application No. GB 2,095,261 (1981); and
French Application No. FR 2,522,655 (1983).

Examples of somatostatin agonists include, but are not limited to, the following somatostatin analogs which are disclosed in the above-cited references:

- 30 H-D- β -Nal-Cys-Tyr-D-Trp-Lys-Thr-Cys-Thr-NH₂ (BIM-23014);

H-D-Phe-Cys-Phe-D-Trp-Lys-Thr-Cys- β -Nal-NH₂;

H-D-Phe-Cys-Tyr-D-Trp-Lys-Thr-Cys- β -Nal-NH₂;

H-D- β -Nal-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

H-D-Phe-Cys-Tyr-D-Trp-Lys-Thr-Pen-Thr-NH₂;

5 H-D-Phe-Cys-Phe-D-Trp-Lys-Thr-Pen-Thr-NH₂;

H-D-Phe-Cys-Tyr-D-Trp-Lys-Thr-Pen-Thr-OH;

H-D-Phe-Cys-Phe-D-Trp-Lys-Thr-Pen-Thr-OH;

H-Gly-Pen-Phe-D-Trp-Lys-Thr-Cys-Thr-OH;

H-Phe-Pen-Tyr-D-Trp-Lys-Thr-Cys-Thr-OH;

10 H-Phe-Pen-Phe-D-Trp-Lys-Thr-Pen-Thr-OH;

H-D-Phe-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-ol (Octreotide);

H-D-Phe-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

H-D-Trp-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;

H-D-Trp-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

15 H-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;

H-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Trp-NH₂;

H-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;

Ac-D-Phe-Lys^{*}-Tyr-D-Trp-Lys-Val-Asp-Thr-NH₂ (an amide
bridge formed between Lys^{*} and Asp);

20 Ac-hArg(Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

Ac-D-hArg(Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

Ac-D-hArg(Bu)-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

Ac-D-hArg(Et)₂-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

Ac-L-hArg(Et)₂-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

25 Ac-D-hArg(CH₂CF₃)₂-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

Ac-D-hArg(CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

Ac-D-hArg(CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Phe-NH₂;

Ac-D-hArg(CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-

NHET;

Ac-L-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

Ac-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys (Me)-Thr-Cys-Thr-NH₂;

5 Ac-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys (Me)-Thr-Cys-Thr-NHEt;

Ac-hArg (CH₃, hexyl)-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

H-hArg (hexyl)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

10 Ac-D-hArg (Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NHEt;

Ac-D-hArg (Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Phe-NH₂;

Propionyl-D-hArg (Et)₂-Gly-Cys-Phe-D-Trp-Lys (iPr)-Thr-Cys-Thr-NH₂;

15 Ac-D-β-Nal-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Gly-hArg (Et)₂-NH₂;

Ac-D-Lys (iPr)-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

Ac-D-hArg (CH₂CF₃)₂-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

20 Ac-D-hArg (CH₂CF₃)₂-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Phe-NH₂;

Ac-D-hArg (Et)₂-D-hArg (Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂;

Ac-Cys-Lys-Asn-4-Cl-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-Ser-D-Cys-NH₂;

25 H-Bmp-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;

H-Bmp-Tyr-D-Trp-Lys-Val-Cys-Phe-NH₂;

H-Bmp-Tyr-D-Trp-Lys-Val-Cys-p-Cl-Phe-NH₂;

H-Bmp-Tyr-D-Trp-Lys-Val-Cys-β-Nal-NH₂;

H-D-β-Nal-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;

30 H-D-Phe-Cys-Tyr-D-Trp-Lys-Abu-Cys-Thr-NH₂;

H-D-Phe-Cys-Tyr-D-Trp-Lys-Abu-Cys- β -Nal-NH₂;
 H-pentafluoro-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂;
 Ac-D- β -Nal-Cys-pentafluoro-Phe-D-Trp-Lys-Val-Cys-Thr-
 NH₂;

- 5 H-D- β -Nal-Cys-Tyr-D-Trp-Lys-Val-Cys- β -Nal-NH₂;
 H-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys- β -Nal-NH₂;
 H-D- β -Nal-Cys-Tyr-D-Trp-Lys-Abu-Cys-Thr-NH₂;
 H-D-p-Cl-Phe-Cys-Tyr-D-Trp-Lys-Abu-Cys-Thr-NH₂;
 Ac-D-p-Cl-Phe-Cys-Tyr-D-Trp-Lys-Abu-Cys-Thr-NH₂;
- 10 H-D-Phe-Cys- β -Nal-D-Trp-Lys-Val-Cys-Thr-NH₂;
 H-D-Phe-Cys-Tyr-D-Trp-Lys-Cys-Thr-NH₂;
 cyclo(Pro-Phe-D-Trp-N-Me-Lys-Thr-Phe);
 cyclo(Pro-Phe-D-Trp-N-Me-Lys-Thr-Phe);
 cyclo(Pro-Phe-D-Trp-Lys-Thr-N-Me-Phe);
- 15 cyclo(N-Me-Ala-Tyr-D-Trp-Lys-Thr-Phe);
 cyclo(Pro-Tyr-D-Trp-Lys-Thr-Phe);
 cyclo(Pro-Phe-D-Trp-Lys-Thr-Phe);
 cyclo(Pro-Phe-L-Trp-Lys-Thr-Phe);
 cyclo(Pro-Phe-D-Trp(F)-Lys-Thr-Phe);
- 20 cyclo(Pro-Phe-Trp(F)-Lys-Thr-Phe);
 cyclo(Pro-Phe-D-Trp-Lys-Ser-Phe);
 cyclo(Pro-Phe-D-Trp-Lys-Thr-p-Cl-Phe);
 cyclo(D-Ala-N-Me-D-Phe-D-Thr-D-Lys-Trp-D-Phe);
 cyclo(D-Ala-N-Me-D-Phe-D-Val-Lys-D-Trp-D-Phe);
- 25 cyclo(D-Ala-N-Me-D-Phe-D-Thr-Lys-D-Trp-D-Phe);
 cyclo(D-Abu-N-Me-D-Phe-D-Val-Lys-D-Trp-D-Tyr);
 cyclo(Pro-Tyr-D-Trp-t-4-AchxAla-Thr-Phe);
 cyclo(Pro-Phe-D-Trp-t-4-AchxAla-Thr-Phe);
 cyclo(N-Me-Ala-Tyr-D-Trp-Lys-Val-Phe);

14


- cyclo(N-Me-Ala-Tyr-D-Trp-t-4-AchxAla-Thr-Phe);
 cyclo(Pro-Tyr-D-Trp-4-Amphe-Thr-Phe);
 cyclo(Pro-Phe-D-Trp-4-Amphe-Thr-Phe);
 cyclo(N-Me-Ala-Tyr-D-Trp-4-Amphe-Thr-Phe);
 5 cyclo(Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba);
 cyclo(Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba-Gaba);
 cyclo(Asn-Phe-D-Trp-Lys-Thr-Phe);
 cyclo(Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-NH(CH₂)₄CO);
 cyclo(Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-β-Ala);
 10 cyclo(Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-D-Glu)-OH;
 cyclo(Phe-Phe-D-Trp-Lys-Thr-Phe);
 cyclo(Phe-Phe-D-Trp-Lys-Thr-Phe-Gly);
 cyclo(Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba);
 cyclo(Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Gly);
 15 cyclo(Asn-Phe-Phe-D-Trp(F)-Lys-Thr-Phe-Gaba);
 cyclo(Asn-Phe-Phe-D-Trp(NO₂)-Lys-Thr-Phe-Gaba);
 cyclo(Asn-Phe-Phe-Trp(Br)-Lys-Thr-Phe-Gaba);
 cyclo(Asn-Phe-Phe-D-Trp-Lys-Thr-Phe(I)-Gaba);
 cyclo(Asn-Phe-Phe-D-Trp-Lys-Thr-Tyr(But)-Gaba);
 20 cyclo(Bmp-Lys-Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-Pro-
 Cys)-OH;
 cyclo(Bmp-Lys-Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-Pro-
 Cys)-OH;
 cyclo(Bmp-Lys-Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-Tpo-
 25 Cys)-OH;
 cyclo(Bmp-Lys-Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-MeLeu-
 Cys)-OH;
 cyclo(Phe-Phe-D-Trp-Lys-Thr-Phe-Phe-Gaba);
 cyclo(Phe-Phe-D-Trp-Lys-Thr-Phe-D-Phe-Gaba);
 30 cyclo(Phe-Phe-D-Trp(5F)-Lys-Thr-Phe-Phe-Gaba);

15

cyclo(Asn-Phe-Phe-D-Trp-Lys(Ac)-Thr-Phe-NH-(CH₂)₃-CO);
 cyclo(Lys-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba);
 cyclo(Lys-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba);
 cyclo(Orn-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba);
 5 H-Cys-Phe-Phe-D-Trp-Lys-Thr-Phe-Cys-NH₂ (BIM-23268);
 H-Cys-Phe-Phe-D-Trp-Lys-Ser-Phe-Cys-NH₂ (BIM-23284);
 H-Cys-Phe-Tyr-D-Trp-Lys-Thr-Phe-Cys-NH₂ (BIM-23295); and
 H-Cys-Phe-Tyr(I)-D-Trp-Lys-Thr-Phe-Cys-NH₂ (BIM-23313).

Note that for all somatostatin agonists described
 10 herein, each amino acid residue represents the structure
 of -NH-C(R)H-CO-, in which R is the side chain (e.g., CH₃
 for Ala) except for Thr-ol which means -NH-CH(CH(CH₃)OH)-
 CH₂-OH and Pro which means prolinyl. Lines between amino
 acid residues represent peptide bonds which join the
 15 amino acids. Also, where the amino acid residue is
 optically active, it is the L-form configuration that is
 intended unless D-form is expressly designated. A
 disulfide bridge is formed between the two free thiols
 (e.g., Cys, Pen, or Bmp residues); however, it is not
 20 shown.

Use of linear somatostatin agonists of the
 following formula is also within the invention:



wherein

A¹ is a D- or L- isomer of Ala, Leu, Ile, Val,
 30 Nle, Thr, Ser, β-Nal, β-Pal, Trp, Phe, 2,4-dichloro-Phe,
 pentafluoro-Phe, p-X-Phe, or o-X-Phe, wherein X is CH₃,
 Cl, Br, F, OH, OCH₃ or NO₂;

16

A² is Ala, Leu, Ile, Val, Nle, Phe, β -Nal, pyridyl-Ala, Trp, 2,4-dichloro-Phe, pentafluoro-Phe, o-X-Phe, or p-X-Phe, wherein X is CH₃, Cl, Br, F, OH, OCH₃ or NO₂;

5 A³ is pyridyl-Ala, Trp, Phe, β -Nal, 2,4-dichloro-Phe, pentafluoro-Phe, o-X-Phe, or p-X-Phe, wherein X is CH₃, Cl, Br, F, OH, OCH₃ or NO₂;

A⁶ is Val, Ala, Leu, Ile, Nle, Thr, Abu, or Ser;

A⁷ is Ala, Leu, Ile, Val, Nle, Phe, β -Nal, 10 pyridyl-Ala, Trp, 2,4-dichloro-Phe, pentafluoro-Phe, o-X-Phe, or p-X-Phe, wherein X is CH₃, Cl, Br, F, OH, OCH₃ or NO₂;

A⁸ is a D- or L-isomer of Ala, Leu, Ile, Val, Nle, Thr, Ser, Phe, β -Nal, pyridyl-Ala, Trp, 2,4-dichloro-Phe, 15 pentafluoro-Phe, p-X-Phe, or o-X-Phe, wherein X is CH₃, Cl, Br, F, OH, OCH₃ or NO₂;

each R₁ and R₂, independently, is H, lower acyl or lower alkyl; and R₃ is OH or NH₂; provided that at least one of A¹ and A⁶ and one of A² and A⁷ must be an aromatic amino acid; and further provided that A¹, A², A⁷ and A⁸ 20 cannot all be aromatic amino acids.





Examples of linear agonists to be used in the method of this invention include:

H-D-Phe-p-chloro-Phe-Tyr-D-Trp-Lys-Thr-Phe-Thr-NH₂;
 25 H-D-Phe-p-NO₂-Phe-Tyr-D-Trp-Lys-Val-Phe-Thr-NH₂;
 H-D-Nal-p-chloro-Phe-Tyr-D-Trp-Lys-Val-Phe-Thr-NH₂;
 H-D-Phe-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-NH₂ (BIM-23052);
 H-D-Phe-Phe-Tyr-D-Trp-Lys-Val-Phe-Thr-NH₂;
 H-D-Phe-p-chloro-Phe-Tyr-D-Trp-Lys-Val-Phe-Thr-NH₂;

30 and

H-D-Phe-Ala-Tyr-D-Trp-Lys-Val-Ala- β -D-Nal-NH₂.

If desired, one or more chemical moieties, e.g., a sugar derivative, mono or poly-hydroxy C₂₋₁₂ alkyl, mono or poly-hydroxy C₂₋₁₂ acyl groups, or a piperazine derivative, can be attached to the somatostatin agonist, e.g., to the N-terminus amino acid. See PCT Application WO 88/02756, European Application 0 329 295, and PCT Application No. WO 94/04752. An example of a somatostatin agonists which contain N-terminal chemical substitutions are:



25 Synthesis of somatostatin agonists

The methods for synthesizing somatostatin agonists is well documented and are within the ability of a person of ordinary skill in the art.

Synthesis of short amino acid sequences is well established in the peptide art. For example, synthesis of H-D-Phe-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-NH₂, described above, can be achieved by following the protocol set forth in Example I of European Patent Application 0 395 417 A1. The synthesis of somatostatin agonists with a substituted N-terminus can be achieved, for example, by following the protocol set forth in WO 88/02756, European Patent Application No. 0 329 295, and PCT Publication No. WO 94/04752.

Somatostatin Receptor Binding Assays

The human SSTR-1, SSTR-2, SSTR-3, SSTR-4, and SSTR-5 cDNA clones have been described (SSTR-1 and SSTR-2 in Yamada, Y., et al., Proc. Natl. Acad. Sci. USA, 89:251-255 (1992); SSTR-3 in Yamada, et al., Mol. Endocrinol. 6:2136-2142 (1993); and SSTR-4 and SSTR-5 in Yamada, et al., Biochem. Biophys. Res. Commun. 195:844-852 (1993)) and are also available from American Type Culture Collection (ATCC, Rockville, MD) (ATCC Nos. 79044 (SSTR-1), 79046 (SSTR-2), and 79048 (SSTR-3)). Based on the restriction endonuclease maps, the entire coding region of each SSTR cDNA may be excised by suitable restriction endonuclease digestion (Maniatis, T., et al., *Molecular Cloning - A Laboratory Manual*, CSHL, 1982). Restriction endonucleases are available from New England Biolabs (Beverly, MA). This cDNA fragment was inserted into the mammalian expression vector, pCMV (Russell, D., et al., J. Biol. Chem., 264:8222-8229 (1989)), using standard molecular biology techniques (see e.g.,

Maniatis, T., et al., Molecular Cloning, -A Laboratory Manual, Cold Spring Harbor Laboratory, 1982) to produce the expression plasmid, pCMV-human SSTR-1 through pCMV-human SSTR-5. Other mammalian expression vectors include
5 pcDNA1/Amp (Invitrogen, Sandlesy, CA). The expression plasmids were introduced into the suitable bacterial host, E. Coli HB101 (Stratagene, La Jolla, CA) and plasmid DNAs, for transfection, were prepared on Cesium Chloride gradients.

10 CHO-K1 (ovary, Chinese hamster) cells were obtained from ATCC (ATCC No. CCL 61). The cells were grown and maintained in Ham's F12 media (Gibco BRL, Grand Island, NY) supplemented with 10% fetal bovine serum under standard tissue culture conditions. For
15 transfection, the cells were seeded at a density 1×10^6 /60-cm plate (Baxter Scientific Products, McGaw Park, IL.). DNA mediated transfection was carried out using the calcium phosphate co-precipitation method (Ausubel, F.M., et al., Current Protocols in Molecular Biology,
20 John Wiley & Sons, 1987). The plasmid pRSV-neo (ATCC; ATCC No. 37198) was included as a selectable marker at 1/10 the concentration of the expression plasmid. CHO-K1 clonal cell lines that have stably inherited the transfected DNA were selected for growth in Ham's F12
25 media containing 10% fetal bovine serum and 0.5mg/ml of G418 (Sigma). The cells were ring-cloned and expanded in the same media for analysis.

Expression of the human SSTR-1 through SSTR-5 receptors in the CHO-K1 cells were detected by Northern
30 blot analysis of total RNA prepared from the cells

(Sambrook, J.E., et al., Molecular Cloning - A Laboratory Manual, Ed. 2., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1989) and by receptor binding using [¹²⁵I-Tyr¹¹]somatostatin-14 as a ligand. Transfected cell
5 lines expressing the human SSTR receptors were clonally expanded in culture and used in the following SSTR binding protocol.

Crude membranes were prepared by homogenization of the transfected cells in 20 ml of ice-cold 50 mM Tris-HCl
10 with a POLYTRON homogenizer (setting 6, 15 sec). Buffer was added to obtain a final volume of 40 ml, and the homogenate was centrifuged in a Sorval SS-34 rotor at 39,000 g for 10 min at 0-4°C. The resulting supernatant was decanted and discarded. The pellet was rehomogenized
15 in ice-cold buffer, diluted, and centrifuged as before. The final pellet was resuspended in the 10 mM Tris HCl and held on ice for the receptor binding assay.

Aliquots of the membrane preparation were
20 incubated for 30 min at 30°C with 0.05 nM [¹²⁵I-Tyr¹¹]somatostatin-14 (2000 Ci/mmol; Amersham Corp., Arlington Heights, IL) in 50 mM HEPES (pH 7.4) containing a test somatostatin agonist of various concentrations (e.g., 10⁻¹¹ to 10⁻⁶), 10 mg/ml bovine serum albumin
25 (fraction V) (Sigma Chemical Co., St. Louis, MO), MgCl₂ (5 mM), Trasylol (200 KIU ml), bacitracin (0.02 mg/ml), and phenylmethylsulphonyl fluoride (0.02 mg/ml). The final assay volume was 0.3 ml. The incubations were terminated by rapid filtration through GF/C filters (pre-soaked in
30 0.3% polyethylenimine for 30 min) using a Brandel

filtration manifold. Each tube and filter were then washed three times with 5 ml aliquots of ice-cold buffer.

Specific binding was defined as the total [125 I-Tyr 11]SRIF-14 bound minus that bound in the presence of 1000 nM. The K_i values for the tested somatostatin agonists were calculated by using the following formula:

$$K_i = IC_{50} / [1 + (LC/LEC)]$$
 where IC_{50} is the concentration of test somatostatin agonist required to inhibit 50 percent of the specific binding of the radioligand [125 I-Tyr 11]somatostatin-14, LC is the concentration of the radioligand (0.05 nM), and LEC is the equilibrium dissociation constant of the radioligand (0.16 nM). The K_i values (nM) for the tested somatostatin agonists are shown in Table I.

TABLE I

	hSSTR-1	hSSTR-2	hSSTR-3	hSSTR-4	hSSTR-5
Somatostatin-14	2.26	0.23	1.2	1.8	1.41
Somatostatin-28	2.38	0.30	1.3	7.93	0.4
Octreotide	875	0.57	26.8	5029	6.78
BIM-23014	2414	0.75	97.9	1826	5.21
BIM-23052	97.6	11.96	5.6	127	1.22
BIM-23190	9120	0.35	215	7537	11.1
BIM-23197	6016	0.19	26.8	3897	9.81
BIM-23272	47.7	3.23	10.9	753	1.01
BIM-23284	27.9	19.3	35.6	58.6	0.85
BIM-23295	86.9	6.19	9.7	3.4	0.34
BIM-23313	15.1	4.78	25.5	55.3	0.30
BIM-26268	1227	15.06	545	3551	0.42

Weight Loss Studies

The effect of chronic (6 day) treatment with BIM-23268 on body weight gain/loss was examined in an obese animal model, the fatty (fa/fa) Zucker rats (purchased from Harlan-Olac, Bicester, Oxon, U.K. See Bray, G., Federation Proceedings 36:148-153 (1977). Eleven male fatty Zucker rats weighing about 450 grams were randomly divided into two groups, and their initial body weights recorded. The animals were housed in pairs in a normal 12 hour light:12 hour darkness cycle at $20 \pm 2^\circ\text{C}$ and fed overnight *ad libitum*.

For the group assigned to receive drug treatment, the rats received the type-5 somatostatin receptor selective agonist BIM-23268C at 3 mg/kg, by subcutaneous injection twice a day at 10:00 a.m. and 5:00 p.m. The other group was treated with a subcutaneous injection of 0.1 ml/100 g of saline twice a day at 10:00 a.m. and 5:00 p.m. The animals were subjected to the BIM-23268 or saline treatment for a total of six days.

At 10:00 a.m. each day, food was removed and replaced with accurately weight 100 gram food pellet (a standard laboratory rat diet, Beekay rat and mouse diet, Bantin & Kingman, Hull, Humberside, U.K.). The amount of food remaining a 10:00 a.m. the next day was accurately weighed, recorded and replaced with 100 grams of fresh food pellets.

The animals were weighed each day during the 6-day treatment period at 5:00 p.m. The untreated control group mean weight was 414.09 at the start of the trial

23

and was 418.89 after six days. The BIM-23268 treated group's mean weight was 413.6 at the start of the trial and remained at 413.6 after six days. The average food consumption for the control group was 26.0 g/rat/day and
5 for the BIM-26268 group was 25.9 g/rat/day.

These results showed that body weight gain was lower in animals treated with BIM-23268. The effect on body weight change was not due to a toxic effect of the agent, as the treated group appeared healthy, and there
10 was no difference in the amount of food consumed over the entire treatment period.

Secondary Endpoints of Efficacy

Because of the amount of weight that must be lost
15 to achieve a clinically important alteration in risk for various sequelae of obesity, the Food and Drug Administration guidelines for the evaluation of weight-control drugs have recommended that additional endpoints showing a decrease in risk factors such as lipemia be
20 monitored.

Obese (fa/fa) Zucker rats were treated as in example 1 above. On the last day of treatment (day 6), food was removed at 5:00 p.m., and the rats were fasted overnight. At 9:00 a.m. the next day, the animals were
25 subjected to a glucose challenge, given as 0.8 gram/kg of glucose orally. Periodic 400 µl of blood samples were taken from the tail vein (Peterson, R.G., ILAR News, 32:16-19 (1990)) 60 min. and 30 min. before and at 30, 60, 90, and 120 min. after the administration of the
30 glucose challenge (0.8 gram/kg orally). Aprotinin

(Traysylol, Bayer UK, Hayward's Health, W. Sussex, U.K.) and heparin (Sigma Chemical Co., Poole, Dorset, U.K.) were added to the blood samples to a final concentration of 400 KIU/ml and 100 units/ml, respectively. Plasma
5 fractions were prepared from these samples by centrifugation at 4000 x G in a microfuge, for the estimation of triglycerides and glycerol. Samples were then stored at -80°C until assayed.

Plasma glycerol and triglycerides were determined
10 using the Sigma Enzymatic (Tinder) calorimetric assay kit (Cat #337-B, Sigma Chemical Co., Poole, Dorset, U.K.) and measuring absorbance at 540 nm in a spectrophotometer.

After six days of treatment with BIM-23268C at 3 mg/kg twice a day by subcutaneous injection, both plasma
15 glycerol and triglycerides were significantly lowered, as exemplified by the samples taken at tim 30 and 60 minutes before the oral glucose challenge. See Fig. 1 and Fig. 2. The administration of an oral glucose challenge have no significant effect on plasma lipids. The BIM-23628C
20 treated group showed a significantly lower plasma glycerol and triglycerides throughout the 2-hour test period. The results suggested that BIM-23268C, following a 6-day treatment period at the prescribed dose was effective in reducing hypertriglyceridemia.

25

Assessment of Efficacy in Patient

The effect of the somatostatin agonist on obesity can be examined in patients by assessing total body weight, body mass index, total adipose tissue content,
30 subcutaneous tissue content, visceral adipose tissue

content (see, e.g., Zamboni, M., Amer. J. Clin. Nutr. 60:682-687 (1994)). The effect of the somatostatin agonist can also be measured on other secondary endpoints, such as insulin sensitivity (see, e.g.,
5 Bergman, R.N., et al., Endocrin. Rev. 6:45-86 (1985); Turner, R.C., Diabetes 44:1-10 (1995)), blood pressure (see, e.g., Maheux, P., Hypertension 24:695-698 (1994)), plasma lipids (see, e.g., Dubrey, S.W., et al., Diabetes 43:831-835 (1994)), and the other acceptable endpoints
10 recommended by the FDA Draft Guidelines for the Clinical Evaluation of Weight Control Drugs (1994) (see also, Drug & Market Development 6:36 (1994)).

OTHER EMBODIMENTS

15 The foregoing description has been limited to specific embodiments of this invention. It will be apparent, however, that variations and modifications may be made to the invention, with the attainment of some or all of the advantages of the invention. Such embodiments
20 are also within the scope of the following claims.

CLAIMS

1. A method of decreasing body weight in a patient, said method comprising administering a therapeutically effective amount of somatostatin or a somatostatin agonist to said patient.
2. A method of claim 1, wherein said method comprises administering a therapeutically effective amount of a somatostatin agonist to said patient.
3. A method of claim 2, wherein said somatostatin agonist is a somatostatin type-2 receptor agonist.
4. A method of claim 2, wherein said somatostatin agonist is a somatostatin type-5 receptor agonist.
5. A method of claim 3, wherein said somatostatin type-2 receptor agonist has a K_i of less than 2 nM for the somatostatin type-2 receptor.
6. A method of claim 4, wherein said somatostatin type-5 receptor agonist has a K_i of less than 2 nM for the somatostatin type-5 receptor.
7. A method of claim 2, wherein said somatostatin agonist is a somatostatin type-2 receptor selective agonist.
8. A method of claim 2, wherein said somatostatin agonist is a somatostatin type-5 receptor selective agonist.
9. A method of claim 7, wherein said somatostatin type-2 receptor selective agonist has a K_i for the somatostatin type-2 receptor that is at least 10 times less than the K_i for the somatostatin type-1, type-3, type-4, and type-5 receptors.

10. A method of claim 8, wherein said somatostatin type-5 receptor selective agonist has a K_i for the somatostatin type-5 receptor that is at least 10 times less than the K_i for the somatostatin type-1, type-2, type-3, and type-4 receptors.
11. A method of decreasing body weight in a patient, said method comprising administering a therapeutically effective amount of H-Cys-Phe-Phe-D-Trp-Lys-Thr-Phe-Cys-NH₂, wherein a disulfide bond exists between the free thiols of two Cys residues.
12. A method of claim 1, wherein said patient is an non-insulin-dependent diabetic human.
13. A method of claim 2, wherein said patient is an non-insulin-dependent diabetic human.
14. A method of claim 3, wherein said patient is an non-insulin-dependent diabetic human.
15. A method of claim 4, wherein said patient is an non-insulin-dependent diabetic human.
16. A method of claim 5, wherein said patient is an non-insulin-dependent diabetic human.
17. A method of claim 6, wherein said patient is an non-insulin-dependent diabetic human.
18. A method of claim 7, wherein said patient is an non-insulin-dependent diabetic human.
19. A method of claim 8, wherein said patient is an non-insulin-dependent diabetic human.
20. A method of claim 9, wherein said patient is an non-insulin-dependent diabetic human.
21. A method of claim 10, wherein said patient is an non-insulin-dependent diabetic human.

22. A method of claim 11, wherein said patient is an non-insulin-dependent diabetic human.

23. A method according to claim 1 wherein the somatostatin agonist is

- 5 H-D- β -Nal-Cys-Tyr-D-Trp-Lys-Thr-Cys-Thr-NH₂,
H-D-Phe-Cys-Phe-D-Trp-Lys-Thr-Cys- β -Nal-NH₂,
H-D-Phe-Cys-Tyr-D-Trp-Lys-Thr-Cys- β -Nal-NH₂,
H-D- β -Nal-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂,
H-D-Phe-Cys-Tyr-D-Trp-Lys-Thr-Pen-Thr-NH₂,
10 H-D-Phe-Cys-Phe-D-Trp-Lys-Thr-Pen-Thr-NH₂,
H-D-Phe-Cys-Tyr-D-Trp-Lys-Thr-Pen-Thr-OH,
H-D-Phe-Cys-Phe-D-Trp-Lys-Thr-Pen-Thr-OH,
H-Gly-Pen-Phe-D-Trp-Lys-Thr-Cys-Thr-OH,
H-Phe-Pen-Tyr-D-Trp-Lys-Thr-Cys-Thr-OH,
15 H-Phe-Pen-Phe-D-Trp-Lys-Thr-Pen-Thr-OH,
H-D-Phe-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-ol
H-D-Phe-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂,
H-D-Trp-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂,
H-D-Trp-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂,
20 H-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂,
H-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Trp-NH₂,
H-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂,
Ac-D-Phe-Lys^{*}-Tyr-D-Trp-Lys-Val-Asp-Thr-NH₂ (an amide
bridge formed between Lys^{*} and Asp),
25 Ac-hArg(Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂,
Ac-D-hArg(Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂,
Ac-D-hArg(Bu)-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂,
Ac-D-hArg(Et)₂-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂,
Ac-L-hArg(Et)₂-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂,
30 Ac-D-hArg(CH₂CF₃)₂-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂,

- Ac-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂,
 Ac-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Phe-NH₂,
 Ac-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NHEt,
 Ac-L-hArg (CH₂-CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂,
 5 Ac-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys (Me) -Thr-Cys-Thr-
 NH₂,
 Ac-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys (Me) -Thr-Cys-Thr-
 NHEt,
 Ac-hArg (CH₃, hexyl) -Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂,
 10 H-hArg (hexyl)₂ -Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂,
 Ac-D-hArg (Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NHEt,
 Ac-D-hArg (Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Phe-NH₂,
 Propionyl-D-hArg (Et)₂-Gly-Cys-Phe-D-Trp-Lys (iPr) -Thr-Cys-
 Thr-NH₂,
 15 Ac-D-β-Nal-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Gly-hArg (Et)₂-
 NH₂,
 Ac-D-Lys (iPr) -Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-Thr-NH₂,
 Ac-D-hArg (CH₂CF₃)₂-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-
 Thr-Cys-Thr-NH₂,
 20 Ac-D-hArg (CH₂CF₃)₂-D-hArg (CH₂CF₃)₂-Gly-Cys-Phe-D-Trp-Lys-
 Thr-Cys-Phe-NH₂,
 Ac-D-hArg (Et)₂-D-hArg (Et)₂-Gly-Cys-Phe-D-Trp-Lys-Thr-Cys-
 Thr-NH₂,
 Ac-Cys-Lys-Asn-4-Cl-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-Ser-D-
 25 Cys-NH₂,
 H-Bmp-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂,
 H-Bmp-Tyr-D-Trp-Lys-Val-Cys-Phe-NH₂,
 H-Bmp-Tyr-D-Trp-Lys-Val-Cys-p-Cl-Phe-NH₂,
 H-Bmp-Tyr-D-Trp-Lys-Val-Cys-β-Nal-NH₂,
 30 H-D-β-Nal-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂,

- H-D-Phe-Cys-Tyr-D-Trp-Lys-Abu-Cys-Thr-NH₂,
H-D-Phe-Cys-Tyr-D-Trp-Lys-Abu-Cys-β-Nal-NH₂,
H-pentafluoro-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-Thr-NH₂,
Ac-D-β-Nal-Cys-pentafluoro-Phe-D-Trp-Lys-Val-Cys-Thr-NH₂,
5 H-D-β-Nal-Cys-Tyr-D-Trp-Lys-Val-Cys-β-Nal-NH₂,
H-D-Phe-Cys-Tyr-D-Trp-Lys-Val-Cys-β-Nal-NH₂,
H-D-β-Nal-Cys-Tyr-D-Trp-Lys-Abu-Cys-Thr-NH₂,
H-D-p-Cl-Phe-Cys-Tyr-D-Trp-Lys-Abu-Cys-Thr-NH₂,
Ac-D-p-Cl-Phe-Cys-Tyr-D-Trp-Lys-Abu-Cys-Thr-NH₂,
10 H-D-Phe-Cys-β-Nal-D-Trp-Lys-Val-Cys-Thr-NH₂,
H-D-Phe-Cys-Tyr-D-Trp-Lys-Cys-Thr-NH₂,
cyclo(Pro-Phe-D-Trp-N-Me-Lys-Thr-Phe),
cyclo(Pro-Phe-D-Trp-N-Me-Lys-Thr-Phe),
cyclo(Pro-Phe-D-Trp-Lys-Thr-N-Me-Phe),
15 cyclo(N-Me-Ala-Tyr-D-Trp-Lys-Thr-Phe),
cyclo(Pro-Tyr-D-Trp-Lys-Thr-Phe),
cyclo(Pro-Phe-D-Trp-Lys-Thr-Phe),
cyclo(Pro-Phe-L-Trp-Lys-Thr-Phe),
cyclo(Pro-Phe-D-Trp(F)-Lys-Thr-Phe),
20 cyclo(Pro-Phe-Trp(F)-Lys-Thr-Phe),
cyclo(Pro-Phe-D-Trp-Lys-Ser-Phe),
cyclo(Pro-Phe-D-Trp-Lys-Thr-p-Cl-Phe),
cyclo(D-Ala-N-Me-D-Phe-D-Thr-D-Lys-Trp-D-Phe),
cyclo(D-Ala-N-Me-D-Phe-D-Val-Lys-D-Trp-D-Phe),
25 cyclo(D-Ala-N-Me-D-Phe-D-Thr-Lys-D-Trp-D-Phe),
cyclo(D-Abu-N-Me-D-Phe-D-Val-Lys-D-Trp-D-Tyr),
cyclo(Pro-Tyr-D-Trp-t-4-AchxAla-Thr-Phe),
cyclo(Pro-Phe-D-Trp-t-4-AchxAla-Thr-Phe),
cyclo(N-Me-Ala-Tyr-D-Trp-Lys-Val-Phe),

- cyclo (N-Me-Ala-Tyr-D-Trp-t-4-AchxAla-Thr-Phe) ,
cyclo (Pro-Tyr-D-Trp-4-Amphe-Thr-Phe) ,
cyclo (Pro-Phe-D-Trp-4-Amphe-Thr-Phe) ,
cyclo (N-Me-Ala-Tyr-D-Trp-4-Amphe-Thr-Phe) ,
5 cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba) ,
cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba-Gaba) ,
cyclo (Asn-Phe-D-Trp-Lys-Thr-Phe) ,
cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-NH (CH₂)₄CO) ,
cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-β-Ala) ,
10 cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-D-Glu) -OH,
cyclo (Phe-Phe-D-Trp-Lys-Thr-Phe) ,
cyclo (Phe-Phe-D-Trp-Lys-Thr-Phe-Gly) ,
cyclo (Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba) ,
cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Gly) ,
15 cyclo (Asn-Phe-Phe-D-Trp (F) -Lys-Thr-Phe-Gaba) ,
cyclo (Asn-Phe-Phe-D-Trp (NO₂) -Lys-Thr-Phe-Gaba) ,
cyclo (Asn-Phe-Phe-Trp (Br) -Lys-Thr-Phe-Gaba) ,
cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Phe (I) -Gaba) ,
cyclo (Asn-Phe-Phe-D-Trp-Lys-Thr-Tyr (But) -Gaba) ,
20 cyclo (Bmp-Lys-Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-Pro-Cys) -
OH,
cyclo (Bmp-Lys-Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-Pro-Cys) -
OH,
cyclo (Bmp-Lys-Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-Tpo-Cys) -
25 OH,
cyclo (Bmp-Lys-Asn-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-MeLeu-
Cys) -OH,
cyclo (Phe-Phe-D-Trp-Lys-Thr-Phe-Phe-Gaba) ,
cyclo (Phe-Phe-D-Trp-Lys-Thr-Phe-D-Phe-Gaba) ,
30 cyclo (Phe-Phe-D-Trp (5F) -Lys-Thr-Phe-Phe-Gaba) ,

32

cyclo(Asn-Phe-Phe-D-Trp-Lys(Ac)-Thr-Phe-NH-(CH₂)₃-CO),

cyclo(Lys-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba),

cyclo(Lys-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba),

cyclo(Orn-Phe-Phe-D-Trp-Lys-Thr-Phe-Gaba),


5 H-Cys-Phe-Phe-D-Trp-Lys-Thr-Phe-Cys-NH₂ ,

H-Cys-Phe-Phe-D-Trp-Lys-Ser-Phe-Cys-NH₂ ,

H-Cys-Phe-Tyr-D-Trp-Lys-Thr-Phe-Cys-NH₂ or

H-Cys-Phe-Tyr(I)-D-Trp-Lys-Thr-Phe-Cys-NH₂ .

24. A method according to claim 1 wherein the
10 somatostatin agonist is



A¹ is a D- or L- isomer of Ala, Leu, Ile, Val,
Nle, Thr, Ser, β-Nal, β-Pal, Trp, Phe, 2,4-dichloro-Phe,
pentafluoro-Phe, p-X-Phe, or o-X-Phe, wherein X is CH₃,
20 Cl, Br, F, OH, OCH₃ or NO₂;

A² is Ala, Leu, Ile, Val, Nle, Phe, β-Nal,
pyridyl-Ala, Trp, 2,4-dichloro-Phe, pentafluoro-Phe, o-X-
Phe, or p-X-Phe, wherein X is CH₃, Cl, Br, F, OH, OCH₃ or
NO₂;

25 A³ is pyridyl-Ala, Trp, Phe, β-Nal, 2,4-dichloro-
Phe, pentafluoro-Phe, o-X-Phe, or p-X-Phe, wherein X is
CH₃, Cl, Br, F, OH, OCH₃ or NO₂;

A⁶ is Val, Ala, Leu, Ile, Nle, Thr, Abu, or Ser;

A⁷ is Ala, Leu, Ile, Val, Nle, Phe, β-Nal,
30 pyridyl-Ala, Trp, 2,4-dichloro-Phe, pentafluoro-Phe, o-X-
Phe, or p-X-Phe, wherein X is CH₃, Cl, Br, F, OH, OCH₃ or
NO₂;

A⁸ is a D- or L-isomer of Ala, Leu, Ile, Val, Nle, Thr, Ser, Phe, β-Nal, pyridyl-Ala, Trp, 2,4-dichloro-Phe, pentafluoro-Phe, p-X-Phe, or o-X-Phe, wherein X is CH₃, Cl, Br, F, OH, OCH₃ or NO₂;

5 each R₁ and R₂, independently, is H, lower acyl or lower alkyl; and R₃ is OH or NH₂; provided that at least one of A¹ and A⁸ and one of A² and A⁷ must be an aromatic amino acid; and further provided that A¹, A², A⁷ and A⁸ cannot all be aromatic amino acids.

10 25. A method according to claim 24 wherein the linear somatostatin agonist is

H-D-Phe-p-chloro-Phe-Tyr-D-Trp-Lys-Thr-Phe-Thr-NH₂,

H-D-Phe-p-NO₂-Phe-Tyr-D-Trp-Lys-Val-Phe-Thr-NH₂,

H-D-Nal-p-chloro-Phe-Tyr-D-Trp-Lys-Val-Phe-Thr-NH₂,



15 H-D-Phe-Phe-Phe-D-Trp-Lys-Thr-Phe-Thr-NH₂,

H-D-Phe-Phe-Tyr-D-Trp-Lys-Val-Phe-Thr-NH₂,


H-D-Phe-p-chloro-Phe-Tyr-D-Trp-Lys-Val-Phe-Thr-NH₂ or

H-D-Phe-Ala-Tyr-D-Trp-Lys-Val-Ala-β-D-Nal-NH₂.


20 26. A method according to claim 1 wherein the somatostatin agonist is



25



or



5 .

27. A method according to claim 1 wherein said patient is obese.

28. A method according to claim 3 wherein said
10 patient is obese.

29. A method according to claim 4 wherein said patient is obese.

30. A method according to claim 7 wherein said patient is obese.

15 31. A method according to claim 8 wherein said patient is obese.

32. A method according to claim 11 wherein said patient is obese.

33. A pharmaceutical or cosmetic composition
20 comprising a therapeutically or cosmetically effective amount of somatostatin; or a somatostatin agonist; or H-Cys-Phe-D-Trp-Lys-Thr-Phe-Cys-NH₂, wherein a disulfide bond exists between the free thiols of the two Cys residues.

25 34. A pharmaceutical composition as claimed in claim 33 having the features identified in any one of claims 3 to 10 and 23 to 26.

35. Use of a somatostatin, or a somatostatin agonist or H-Cys-Phe-Phe-D-Trp-Lys-Thr-Phe-Cys-NH₂, wherein a disulfide bond exists between the free thiols of the two Cys residues, in the formulation of a
5 pharmaceutical or cosmetic composition for use in reducing excessive body weight in a human or mammalian animal.

36. Use of a somatostatin, or a somatostatin agonist according to claim 35, wherein said somatostatin
10 or somatostatin agonist has the relevant features identified in any one of claims 3 to 10 and 23 to 26.

37. A pharmaceutical composition substantially as hereinbefore described with reference to the Examples.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/02999

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K38/31 A61K7/48

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 11962 A (BIOMEASURE INC ;UNIV TULANE (US); COY DAVID HOWARD (US); TAYLOR JO) 3 April 1997 see page 1, line 1 - line 29 see page 3 - page 4 see page 6, line 12 - line 23 see page 7, line 30 - line 34 ---	33-37
X	CARRETTA R ET AL: "REDUCTION OF BLOOD PRESSURE IN OBESE HYPERINSULINAEMIC HYPERTENSIVE PATIENTS DURING SOMATOSTATIN INFUSION" JOURNAL OF HYPERTENSION, vol. 7, no. SUPPL. 06, 18 June 1989, page S196/S197 XP002053034 see the whole document ----- -/--	33, 34, 37

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

21 September 1998

Date of mailing of the international search report

30/09/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Fernandez y Branas, F

INTERNATIONAL SEARCH REPORT

Inte. Jonal Application No

PCT/EP 98/02999

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	WO 98 10786 A (COHEN YAROM) 19 March 1998 see the whole document ----	1-37
X,P	WO 98 09991 A (UNIV WASHINGTON ;ZYMOGENETICS INC (US)) 12 March 1998 see page 8, line 33 - line 12 see page 30, line 19 - line 25 ----	1,2,12, 13,27, 33,35,37
X	WO 96 35950 A (UNIV BUCKINGHAM) 14 November 1996 see the whole document ----	33,34,37
X	EP 0 657 174 A (MAYO FOUNDATION) 14 June 1995 see the whole document -----	33,34,37

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 98/ 02999

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 1 - 32
are directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/02999

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9711962 A	03-04-1997	US 5708135 A AU 6914596 A EP 0859785 A NO 981395 A	13-01-1998 17-04-1997 26-08-1998 27-03-1998
WO 9810786 A	19-03-1998	AU 4133997 A	02-04-1998
WO 9809991 A	12-03-1998	AU 4250397 A	26-03-1998
WO 9635950 A	14-11-1996	US 5763200 A AU 694360 B AU 5764596 A CA 2220106 A EP 0829011 A	09-06-1998 16-07-1998 29-11-1996 14-11-1996 18-03-1998
EP 0657174 A	14-06-1995	US 5583104 A	10-12-1996